



CORTEVA™
agriscience

syngenta

FMC



BASF
We create chemistry

**CENTRAL
ONTARIO** **FS**

Ontario Crops Research Centre – Bradford (Muck Station) IPM Report 2025

This is the Muck Crops IPM Update for August 11, 2025

Highlights

- **Stemphylium leaf blight severity has really increased in many onion fields and lesions are sporulating**
- Tip dieback is very prominent
- Some seeded onion fields are lodging
- Thrips populations have been increasing in some seeded onion fields as thrips leave nearby lodging onion fields
- Carrot canopies are closing in
- Aster yellows has been found in a number of fields and Aster leafhopper populations are high in some fields
- Some carrot rust fly activity has been observed

Disease Forecasting Highlight:

Botrytis leaf blight (BOTcast)	Onion downy mildew (DOWNcast)	Stemphylium leaf blight (BSPcast)	White Rot on onions	Sclerotinia (White Mold) on Carrot	General conditions for disease (TOMcast)	Lettuce downy mildew (BREMcast)
LOW RISK	LOW RISK	HIGH RISK	LOW-MODERATE RISK	LOW RISK	HIGH RISK	MODERATE RISK

Onion Update

A number of seeded onion fields are starting to lodge. Transplant onions are being harvested.

Stemphylium leaf blight severity has been increasing and many lesions are now sporulating. The number of Stemphylium spores caught in our spore traps has also been increasing. Tip dieback has become more common and severe in some fields, which can be an entry point for further Stemphylium development. If your onions will be standing for a while we recommend alternating the group 7 fungicides (Miravis Duo, Sercadis, Aprovia/Aprovia Top and Merivon) with Allegro (group 29) for Stemphylium leaf blight suppression and resistance management. Resistance has been developing to the active ingredient in Sercadis and Merivon. The group 7



**Sporulating
Stemphylium lesions**
Scout – Dylan Quanbury



ENZA ZADEN



**HOLMES
AGRO**

Ontario



NORSECO



systemic products are effective for around 10-14 days and the contact fungicide Allegro is effective for around 7- 10 days.

We do not recommend spraying Quadris Top (group 11/3) or Luna Tranquility (group 7/9) as Stemphylium has developed resistance to these fungicides. There is very high resistance in Stemphylium to one of the fungicides in Quadris Top and high resistance to one of the fungicides in Luna Tranquility.

Check out our IPM video on Stemphylium leaf blight in the Holland Marsh for more info by clicking on the thumbnail to the right.



Stemphylium Leaf Blight in the Holland Marsh



How to Control Onion Thrips

Thrips populations vary from field to field. If your field surpasses the 3 thrips/leaf threshold the insecticides we would recommend applying next are Agri-Mek SC (group 6), Delegate WG (group 5), Minecto Pro (group 6/28) or Exirel (group 28). These insecticides should also be applied back-to-back, no more than 2 weeks apart. Check out our IPM video on onion thrips for more info by clicking on the thumbnail to the left.

In terms of onion downy mildew, although there is usually a high risk this time of year conditions have not been conducive for the disease this year as it has been too hot and dry.

Scouts are also assessing onion fields for 2nd generation onion maggot damage.

Registered onion herbicides can be found in the [Ontario Crop Protection Hub](#).

Carrot Update

Carrot canopies continue to close in.

Aster yellows has been found in a number of fields. Aster leafhoppers continue to be active and populations are high in some fields. The percent of aster leafhoppers with aster yellows coming from the south has also been fairly high. Sevin XLR, Admire 240F, Actara 25WG, Closer and Sivanto Prime are registered if your field is over the 20 leafhoppers/trap or 1 leafhopper/sweep thresholds.



Aster yellows symptoms
Scout – Dylan Quanbury

Some carrot rust fly activity has been observed. This generation of rust fly can cause damage and it is important to spray if your field is over threshold. If your field is over the 0.1 rust flies/trap/day threshold, or remaining close to it over a long period of time, Mako, UP-Cyde 2.5 EC, Labamba, Matador 120EC and Silencer 120 EC are registered for control.



ENZA ZADEN





CORTEVA
agriscience

syngenta

FMC



BASF
We create chemistry

**CENTRAL
ONTARIO** **FS**

For fields with canopies starting to close in, it may be important to apply a Sclerotinia white mold fungicide soon to get the product down to the soil while you can. Allegro 500F, Scholar 230SC, Luna Sensation and Switch 62.5 WG are registered for control and Miravis Duo and Fontelis are registered for suppression. This may be even more important in high white mold risk fields.

Registered carrot herbicides can be found in the [Ontario Crop Protection Hub](#).

Celery Update

Celery harvest is ongoing. Disease and insect damage has been minimal. Blights have been found regularly in celery, but severity is still low. Celery leaf curl has been very infrequent. Tarnished plant bugs have been active, but damage has been minimal as well.

Registered celery herbicides can be found in the [Ontario Crop Protection Hub](#).

Daily Weather

Date (August)	Max temperature (°C)	Min temperature (°C)	Rain (mm)
7	29.3	16.6	0.0
8	29.5	15.4	0.0
9	33.5	16.1	0.0
10	34.2	17.8	0.0

Soil Temperature (°C): 5cm: 24.0

Any questions or comments? Please call Tyler Blauel or Mary Ruth McDonald at 905-775-3783



**Alliance
Agri-Turf**
Knowledge to grow by



STOKES
SEEDS

ENZA ZADEN



**HOLMES
AGRO**

Ontario



NORSECO