



CORTEVA
agriscience

syngenta

FMC



BASF
We create chemistry

**CENTRAL
ONTARIO** **FS**

Ontario Crops Research Centre – Bradford (Muck Station) IPM Update 2026

Date: August 24

Highlights

- Many onion fields are at various stages of lodging, windrowing and harvesting.
- **For fields with onions that will be standing for a while longer:**
 - o **Onion downy mildew has been confirmed in the Holland Marsh. There continues to be a high risk for onion downy mildew infection with the sporulation infection periods last week.**
 - o **Stemphylium leaf blight symptoms continue to spread. There is a high risk for further disease development.**
- 2nd generation carrot rust flies are active and some fields are over threshold.
- Aster leafhopper activity is sporadic.
- Sclerotinia white mold has been found in several carrot fields.
- The [2026 Muck Crops Research Station Grower Field Day](#) will be held next week on Thursday September 3rd from 8:30am – 4 pm. Coffee and donuts will be served compliments of Stokes Seeds. Lunch will also be provided. We look forward to seeing everyone here.

Disease Forecasting Highlight:

Botrytis leaf blight (BOTcast)	Onion downy mildew (DOWNcast)	Stemphylium leaf blight (BSPcast)	White Rot on onions	Sclerotinia (White Mold) on Carrot	General conditions for disease (TOMcast)	Lettuce downy mildew (BREMcast)
LOW RISK	HIGH RISK	HIGH RISK	LOW RISK	MODERATE RISK	HIGH RISK	HIGH RISK

Onion Update

Onion downy mildew has been found in the Holland Marsh. There continues to be a high risk for onion downy mildew infection with the sporulation infection periods last week. If your onions will be standing for a while longer there are many products registered for control of downy mildew, but our research at the Muck Station has shown that Orondis Ultra (group 40/49), Orondis Gold (group 4/+49) and



HOLMES
AGRO

ENZA ZADEN

STOKES
SEEDS



**Alliance
Agri-Turf**
Knowledge to grow by



NORSECO

SEMINOVA



Ontario






Zampro (group 45/40) are the most effective for controlling this disease. Ridomil Gold MZ (group 4/M3) is no longer being produced, but if you have some it is an effective product as well. Orondis Ultra should be applied with a non-ionic surfactant. These systemic products should protect onions for 10 days. Mancozeb products, such as Manzate Pro-Stick, Dithane Rainshield and Penncozeb 75 DF Raincoat (all group M3), can be effective for controlling downy mildew if applied before infection takes place. Rotating fungicides is very important.

Stemphylium leaf blight is present in all onion fields and disease symptoms have been increasing gradually. Disease severity ranges from field to field; some fields have more severe disease development. More Stemphylium sporulation has been found on onion tips and lower onion lesions. There continues to be a high risk for disease development. We recommend alternating the group 7 fungicides (Miravis Duo, Sercadis, Aprovia/Aprovia Top and Merivon) with Allegro (group 29) for Stemphylium leaf blight suppression and resistance management. Resistance has been developing to the active ingredient in Sercadis and Merivon. The group 7 systemic products are effective for around 10-14 days and the contact fungicide Allegro is effective for around 7- 10 days.

We do not recommend spraying Quadris Top (group 11/3) or Luna Tranquility (group 7/9) for Stemphylium as resistance has developed to these fungicides. Latest research has found very high resistance in Stemphylium to one of the fungicides in Quadris Top and high resistance to one of the fungicides in Luna Tranquility.



Stemphylium symptoms - Scout

Royal MH can be applied to lodging onions to control sprouting in storage. Royal MH should be applied when bulbs are mature, have 5-8 green leaves and ~50% of tops have fallen. Green tops are needed for the product to be absorbed into the bulb. Royal MH should be applied no later than 10 days before harvest.

Over the next two weeks scouts will be completing their final harvest assessment on lodged onion fields.

Carrot Update



White mold - Scout

Sclerotinia white mold has been found in a number of fields. Allegro 500F, Scholar 230SC, Luna Sensation and Switch 62.5 WG are registered for control and Miravis Duo and Fontelis are registered for suppression. It is important to try and get these products down into the crown and lower canopy as best you can. For fields where canopies are closing in it is important to apply a fungicide soon to get the product down to the soil while you can.



ENZA ZADEN



NORSECO



Ontario





CORTEVA
agriscience

syngenta

FMC



BASF
We create chemistry

**CENTRAL
ONTARIO** **FS**

Second generation carrot rust flies are active. This generation can cause damage and some fields are over threshold. If your field is over the 0.1 flies/trap/day threshold Mako, UP-Cyde 2.5 EC, Labamba, Matador 120EC and Silencer 120 EC are registered for control.

Aster leafhopper activity has been variable and aster yellows has been minimal. Sevin XLR, Admire 240F, Actara 25WG, Closer and Sivanto Prime are registered if your field is over the 20 leafhoppers/trap or 1 leafhopper/sweep thresholds.

For weed control, registered [preemergence](#) and [postemergence](#) herbicides can be found in the Ontario Crop Protection Hub by following the links.

Celery Update

Celery harvest continues. Blights have been found regularly, but overall disease severity continues to be low. There are many fungicides registered to manage blights in celery, which can be found on the [Ontario Crop Protection Hub](#). Celery leaf curl has been found in these wet conditions as well. Switch 62.5 WG and Quadris Flowable are registered for leaf curl control. Some pink rot has also been found by scouts. Registered fungicides for pink rot can be found [here](#). Tarnished plant bugs have been active, but damage continues to be minimal.

Registered celery herbicides can be found in the [Ontario Crop Protection Hub](#).

Daily Weather

Date (August)	Max temperature (°C)	Min temperature (°C)	Rain (mm)
20	25.6	9.4	0.0
21	26.3	9.3	0.0
22	26.1	16.1	9.8
23	20.2	13.9	12.6

Soil Temperature (°C): 5cm: 18.3

Any questions or comments? Please call Tyler Blauel at (365) 340-4158



HOLMES
AGRO

STOKES
SEEDS

ENZA ZADEN



Alliance
Agri-Turf
Knowledge to grow by



NORSECO



SEMINOVA



Ontario

