



# Ontario Crops Research Centre – Bradford (Muck Station) IPM Report 2025

This is the Muck Crops IPM Update for June 12, 2025

## Highlights

- Onions are around the first to third true leaf stage
- Scouts are just starting to find thrips, populations are very low and slowly growing
- Most carrot fields are seeded now
- Carrot weevils are active and in their egg laying period; activity is field specific
- 1<sup>st</sup> generation carrot rust flies are active, but this generation cannot cause serious damage

## Disease Forecasting Highlight:

Botrytis leaf blight (BOTcast)	Onion downy mildew (DOWNcast)	Stemphylium leaf blight (BSPcast)	White Rot on onions	Sclerotinia (White Mold) on Carrot	General conditions for disease (TOMcast)	Lettuce downy mildew (BREMcast)
LOW RISK	LOW RISK	LOW RISK	MODERATE RISK	LOW RISK	MODERATE RISK	HIGH RISK

## Onion Update

Onions are generally at the 1<sup>st</sup> to 3<sup>rd</sup> true leaf stage. Transplants are around the 4<sup>th</sup> to 5<sup>th</sup> true leaf stage.

The scouts are starting to find very low numbers of thrips in onion fields, mainly transplants. We will continue to monitor population growth. Populations of thrips need to build more before spraying Movento.

Onion maggot flies activity continues to increase. We will continue to monitor populations throughout the marsh. Seeds treated with Sepresto should be protected against onion maggot damage. Transplant onions have a higher risk of onion maggot damage.

Early season weed control is important. Some goal damage has been observed on transplant and seeded onions. Pre weed emergence herbicides include Chateau, Prowl H2O and Frontier Max. Post weed emergence herbicides include Goal 2XL, Pardner and bromoxynil products. Allowing the onions waxy leaf layer to develop can help protect the plant from herbicide injury. Registered herbicides can be found in the [Ontario Crop Protection Hub](#).



ENZA ZADEN





## Carrot Update

Many carrots are emerging or at the bunny ear stage. Earlier carrots are around the 1<sup>st</sup> to 2<sup>nd</sup> true leaf stage. Carrot weevils are active and are in their egg laying period. Carrot weevil pressure is usually higher in early seeded carrots than later seeded carrots. Carrot weevil populations look to be fairly low overall, but some fields have higher weevil populations. If you have earlier seeded carrots, we recommend spraying for weevils at the 2<sup>nd</sup> true leaf stage if your field is over the cumulative 1.5 weevils/trap threshold. If your field is over the 5 weevils/trap threshold, we also recommend spraying at the 4<sup>th</sup> true leaf stage as well. Rimon 10 EC or Exirel are good insecticides to manage weevil populations.



How to Control Carrot Weevil

For more information on controlling carrot weevils, check out our “How to Control Carrot Weevil” video on our Muck Crops IPM YouTube channel by clicking on the thumbnail to the left.

First generation carrot rust flies are active now as well, but this generation cannot cause serious damage as the carrots are too small at this time.

## Daily Weather

Date (June)	Max temperature (°C)	Min temperature (°C)	Rain (mm)
9	23.0	15.4	0.0
10	19.8	12.2	0.2
11	27.4	14.1	0.0

Soil Temperature (°C): 5cm: 19.2

Any questions or comments? Please call Tyler Blauel or Mary Ruth McDonald at 905-775-3783



ENZA ZADEN

