



IMPROVE LIFE.



Ontario Crops Research Centre – Bradford (Muck Station) IPM Report 2024

This is the Muck Crops IPM Update for June 24, 2024

Highlights

- Conditions have been favourable for Stemphylium leaf blight. Stemphylium has been identified on seeded and transplant onions in the marsh.
- We received ~48 mm of rain since Thursday.
- Most seeded onions are at the 3rd - 4th true leaf stage.
- Thrips have been found in many fields and populations are slowly building.
- Many carrot fields are at the 2nd - 3rd true leaf stage, some are at bunny ears.
- Carrot weevils have almost completed their egg laying period. Some fields are over the 1.5 and 5 cumulative weevils/trap threshold.
- The AgRobotics Working Group will be holding in-field robot demonstrations on July 4th. If you are interested, please register for this free tour [here](#). The tour will start at the Muck Station at 9am.

Disease Forecasting Highlight:

Botrytis leaf blight (BOTcast)	Onion downy mildew (DOWNcast)	Stemphylium leaf blight (BSPcast)	White Rot on onions	Sclerotinia (White Mold) on Carrot	General conditions for disease (TOMcast)	Lettuce downy mildew (BREMcast)
LOW RISK	LOW RISK	HIGH RISK	LOW RISK	LOW RISK	HIGH RISK	LOW RISK

Onion Update

Most seeded onions are around the 3rd-4th true leaf stage. The risk for Stemphylium is currently high after the wet weather we had this past weekend. Stemphylium leaf blight has now been found in seeded onions, as well as transplants.

First spray: If your seed was treated with EverGol Prime or Pen 240 your first spray should be Allegro 500F. Allegro is effective for around 7-10 days. If your seeds were not treated with penflufen, Miravis Duo (Group 7/3) would be the best choice to protect your onions.

Second spray: If it has been 7-10 days since you applied Allegro, the next fungicide you apply could be Miravis Duo (Group 7/3). Aprovia/Aprovia Top, Sercadis and Merivon are also group 7s that are registered to suppress Stemphylium, but resistance has been developing to the active ingredient in Sercadis and Merivon. Rotating these products is important even though they are all group 7s. Allegro is low risk for developing resistance and can be used in rotation with these products.





IMPROVE LIFE.



We do not recommend spraying Quadris Top (group 11/3) or Luna Tranquility (group 7/9) as Stemphylium has developed resistance to these fungicides. There is very high resistance in Stemphylium to one of the fungicides in Quadris Top and high resistance to one of the fungicides in Luna Tranquility.

Some onion maggot damage has been seen in seeded and transplant onions. Damage from 1st generation onion maggots will be assessed.



Madison Sirois - Scout

Thrips populations are slowly building in many fields. Generally, thrips populations are slightly higher in transplant fields compared to seeded fields. For thrips management, it is important to first apply Movento 240 SC once thrips are present in your field and their population is building. Movento should be sprayed back-to-back no more than 2 weeks apart for resistance management.



Eriel Strauch - Scout

For weed control, post weed emergence herbicides include Goal 2XL, Pardner and bromoxynil products. Registered herbicides can be found in the [Ontario Crop Protection Hub](#).

Carrot Update

Most carrot fields are at the 2nd - 3rd true leaf stage. Earlier seeded carrots are around the 4th - 5th true leaf stage. We have seen some heat canker in small carrot fields after the hot weather last week.

The egg laying period for carrot weevils is almost complete. Carrot weevil pressure is usually higher in early seeded carrots than later seeded carrots as later seeded carrots often miss the majority of this egg laying period. If you have earlier seeded carrots, we recommend spraying for weevils at the 2nd true leaf stage if your field is over the cumulative 1.5 weevils/trap threshold. If your field is over the 5 weevils/trap threshold, we also recommend spraying at the 4th true leaf stage. Rimon 10 EC or Exirel are good insecticides to manage weevil populations.





IMPROVE LIFE.



For weed control, registered herbicides can be found in the [Ontario Crop Protection Hub](#).

Celery Update

Some early celery is now being harvested while transplanting is finishing up in other fields. Diseases and insect damage have been minimal in celery so far. Some herbicide damage on the leaves has been common.

Daily Weather

Date (June)	Max temperature (°C)	Min temperature (°C)	Rain (mm)
20	30.2	20.7	9.4
21	25.7	19.8	23.6
22	24.8	19.2	14.4
23	25.7	17.3	0.2

Soil Temperature (°C): 5cm: 21.2

Any questions or comments? Please call Tyler Blauel or Mary Ruth McDonald at 905-775-3783

