

Ontario Crops Research Centre – Bradford (Muck Station) IPM Report 2024

This is the Muck Crops IPM Update for May 28, 2024

Highlights

- We received over 30 mm of rain yesterday.
- Most onion fields have now been seeded, some carrot fields still need to be seeded.
- Onion maggot fly activity continues.
- Carrot weevils continue to be active.
- 1st generation carrot rust flies are active, this generation cannot cause serious damage.

Botrytis leaf	Onion downy	Stemphylium	White Rot	Sclerotinia	General	Lettuce
blight	mildew	leaf blight	on onions	(White Mold)	conditions for	downy
(BOTcast)	(DOWNcast)	(BSPcast)		on Carrot	disease	mildew
					(TOMcast)	(BREMcast)
LOW RISK	LOW RISK	LOW RISK	MODERATE	LOW RISK	MODERATE RISK	HIGH RISK
			RISK			

Disease Forecasting Highlight:

Onion Update

Most onion fields have now been seeded. Many fields are around the 1st true leaf stage. We will continue to monitor onion maggot populations throughout the marsh. Seeds treated with Sepresto should be protected against onion maggot damage. The barley wind abatement has established well in many fields and has been sprayed down in several fields as well.

Early season weed control is important. Pre weed emergence herbicides include Chateau, Prowl H2O and Frontier Max. Post weed emergence herbicides include Goal 2XL, Pardner and bromoxynil products. Allowing the onions waxy leaf layer to develop can help protect the plant from herbicide injury. Registered herbicides can be found in the <u>Ontario Crop Protection Hub</u>.

Carrot Update

Many carrot fields have now been seeded. Carrot weevils are active and are in their egg laying period. Carrot weevil pressure is usually higher in early seeded carrots than later seeded carrots. Some of the earliest seeded carrots in the marsh are around the 1st true leaf stage. We do not recommend spraying for weevils until the 2nd true leaf stage if you are over the cumulative 1.5 weevils/trap threshold. Rimon 10 EC or Exirel are good insecticides to manage weevil populations.







We create chemistry

MINISTRY OF AGRICULTURE, FOOD AND RURAL AFFAIRS



Although 1st generation carrot rust flies are now active, this generation cannot cause serious damage as the carrots are too small or have not emerged at this time.

Daily Weather

Date (May)	Max temperature (°C)	Min temperature (°C)	Rain (mm)
23	24.5	13.5	0.0
24	22.7	11.2	0.0
25	18.7	8.5	3.0
26	23.0	11.2	0.0
27	23.7	14.5	36.6

Soil Temperature (°C): 5cm: 16.6

Any questions or comments? Please call Tyler Blauel or Mary Ruth McDonald at 905-775-3783





